



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification: C12N 5/06, A61P 37/00, C12N 5/08	A2	(11) International Publication Number: WO 00/36089 (43) International Publication Date: 22 June 2000 (22.06.2000)
(21) International Application Number: PCT/GB99/04233 (22) International Filing Date: 15 December 1999 (15.12.1999) (30) Priority Data: 9827604.1 15 December 1998 (15.12.1998) GB (60) Parent Application or Grant LORANTIS LIMITED [/]; (). LAMB, Jonathan, Robert [/]; (). DALLMAN, Margaret, Jane [/]; (). HOYNE, Gerard, Francis [/]; (). LAMB, Jonathan, Robert [/]; (). DALLMAN, Margaret, Jane [/]; (). HOYNE, Gerard, Francis [/]; (). MALLALIEU, Catherine, Louise ; ().	Published	
(54) Title: METHODS OF IMMUNOSUPPRESSION (54) Titre: PROCEDE D'IMMUNOSUPPRESSIONN		
(57) Abstract		
<p>A method for producing a T cell having tolerance to an allergen or antigen which method comprises incubating the T cell with an antigen presenting cell (APC) in the presence of (i) a composition capable of upregulating expression of an endogenous Notch ligand in the APC and (ii) the allergen or antigen is provided.</p>		
(57) Abrégé		
<p>L'invention concerne un procédé de production d'un lymphocyte T tolérant à un allergène ou antigène, qui consiste à incuber le lymphocyte T avec une cellule présentatrice de l'antigène (CPA), en présence i) d'une composition capable de réguler positivement l'expression d'un ligand Notch endogène dans la CPA, et ii) de l'allergène ou antigène.</p>		

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁷ : C12N 5/06, 5/08, A61P 37/00</p>	<p>A2</p>	<p>(11) International Publication Number: WO 00/36089 (43) International Publication Date: 22 June 2000 (22.06.00)</p>
<p>(21) International Application Number: PCT/GB99/04233 (22) International Filing Date: 15 December 1999 (15.12.99) (30) Priority Data: 9827604.1 15 December 1998 (15.12.98) GB (71) Applicant (for all designated States except US): LORANTIS LIMITED [GB/GB]; 38 Jermyn Street, London SW1Y 6DN (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): LAMB, Jonathan, Robert [GB/GB]; 7 Mansion House, Edinburgh EH19 1TZ (GB). DALLMAN, Margaret, Jane [GB/GB]; 64 Grafton Road, Acton, London W3 6PF (GB). HOYNE, Gerard, Francis [AU/GB]; 15 Clayknowes Way, Musselburgh, Midlothian EH21 6UL (GB). (74) Agent: MALLALIEU, Catherine, Louise; D. Young & Co., 21 New Fetter Lane, London EC4A 1DA (GB).</p>		<p>(81) Designated States: AF, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>Without international search report and to be republished upon receipt of that report.</i></p>
<p>(54) Title: METHODS OF IMMUNOSUPPRESSION (57) Abstract <p>A method for producing a T cell having tolerance to an allergen or antigen which method comprises incubating the T cell with an antigen presenting cell (APC) in the presence of (i) a composition capable of upregulating expression of an endogenous Notch ligand in the APC and (ii) the allergen or antigen is provided.</p></p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakistan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Description

5

10

15

20

25

30

35

40

45

50

55

5

METHODS OF IMMUNOSUPPRESSIONField of the Invention

10

5 The present invention relates to methods for preparing antigen presenting cells and lymphocytes, particularly but not exclusively regulatory T cells, that can suppress the activity of lymphocytes and other cells of the immune system. It also relates to the use of compositions capable of upregulating expression of an endogenous Notch or Notch ligand in such methods. These compositions, antigen presenting cells and lymphocytes may be used in immunotherapy.

20

Background to the Invention

25

15 Immunological tolerance to self-antigens is vital to the proper functioning of the mammalian immune system. In addition to the deletion of self-reacting T cells in the thymus, active suppression mediated by regulatory T cells has recently been identified as an important mechanism for maintaining peripheral tolerance (WO98/20142). In autoimmune diseases such as multiple sclerosis, rheumatoid arthritis or diabetes, there is a failure of the proper regulation of tolerance. Improved treatment methods for re-

20 establishing tolerance are desirable for autoimmune diseases. Similarly in allergic conditions and for transplantation of an organ or tissue from a donor individual, induction of tolerance to particular foreign antigens or profiles of foreign antigens is desirable.

35

40

25 It has recently been shown that it is possible to generate a class of regulatory T cells which are able to transmit antigen-specific tolerance to other T cells, a process termed infectious tolerance (WO 98/20142). The functional activity of these cells can be mimicked by over-expression of a Notch ligand protein on their cell surfaces. In particular, regulatory T cells can be generated by over-expression of a member of the Delta or Serrate family of Notch ligand proteins. Delta or Serrate expressing T cells specific to one antigenic epitope are

30 also able to transfer tolerance to T cells recognising other epitopes on the same or related antigens, a phenomenon termed "epitope spreading".

45

50

55

5 WO98/20142 describes methods for generating regulatory T cells by either transfecting
hybridoma T cells with a nucleic acid construct directing the expression of Delta or by
transfecting antigen presenting cells such as dendritic cells with a nucleic acid construct
10 directing the expression of Serrate and incubating the dendritic cells with T cells.

5

Summary of the invention

15 The present invention identifies substances capable of upregulating expression of the
endogenous genes encoding Notch or Notch ligands in antigen presenting cells (APCs) and
10 lymphocytes. We believe that incubating APCs and lymphocytes, e.g. T cells, in the
presence of these substances and a specific antigen produces APCs capable of inducing
20 immunological tolerance in such lymphocytes or other APCs to the specific antigen.
Furthermore, we believe that administration of these APCs and/or lymphocytes to a
recipient individual may induce immunotolerance in that individual to the antigen. In
25 particular we believe that immunosuppressive cytokines (such as IL-4, IL-10, IL-13, TGF-
β and SLIP3 ligand) can be used to upregulate the expression of endogenous Notch or
Notch ligands in APCs or lymphocytes. The present invention applies these findings to the
30 generation of primed APCs and lymphocytes, e.g. regulatory T cells, using *ex vivo*
methods. The resulting primed APCs and/or lymphocytes, e.g. regulatory T cells, may be
20 readministered to the patient to treat or prevent a range of immune disorders resulting from
inappropriate lymphocyte activity, such as auto-immune disease and graft rejection.

35 Accordingly the present invention provides a method for producing a lymphocyte or antigen
presenting cell (APC) having tolerance to an allergen or antigen which method comprises
40 incubating a lymphocyte or APC obtained from a human or animal patient with (i) a
composition capable of upregulating expression of an endogenous Notch or Notch ligand in
the lymphocyte and/or APC and (ii) the allergen or antigen.

45 According to a preferred embodiment the method comprises incubating a lymphocyte or APC
obtained from a human or animal patient with an APC in presence of (i) a composition
capable of upregulating expression of an endogenous Notch or Notch ligand in the
50 lymphocyte and/or APC and (ii) the allergen or antigen.

55

5 Preferably the lymphocyte is a T cell or a B cell. Most preferably the lymphocyte is a T cell.

10 Accordingly the present invention further provides a method for producing *ex vivo* a T cell having tolerance to an allergen or antigen which method comprises incubating a T cell obtained from a human or animal patient with an antigen presenting cell (APC) in the presence of (i) a composition capable of upregulating expression of an endogenous Notch and/or Notch ligand in the APC and/or T cell and (ii) the allergen or antigen.

15 Preferably, the composition comprises a polypeptide selected from Noggin, Chordin, Follistatin, Xnr3, FGF and derivatives, fragments, variants and homologues thereof, and immunosuppressive cytokines, or a combination thereof. More preferably, the composition
20 comprises at least one polypeptide selected from Noggin, Chordin, Follistatin, Xnr3, fibroblast growth factors and derivatives, fragments, variants and homologues thereof, and/or at least one immunosuppressive cytokine. Particularly preferred examples of
25 immunosuppressive cytokines for use in the present invention are IL-4, IL-10, IL-13, TGF- β and FLT3 ligand.

30 The Notch ligand is preferably selected from Serrate, Delta and homologues thereof, more preferably Serrate and Delta.

35 20 The present invention also provides a second method for producing *ex vivo* a lymphocyte or APC having tolerance to an allergen or antigen which method comprises incubating a lymphocyte or APC obtained from a human or animal patient with a lymphocyte or APC produced by the above methods of the invention.

40 25 APCs or lymphocytes, preferably T cells, produced *ex vivo* by the methods of the invention may be used in suppressing an immune response in a mammal to the allergen or antigen, such as an auto-immune response or allograft rejection.

45 We have identified substances and combinations of substances capable of upregulating
30 expression of an endogenous Notch or Notch ligand in an APC or lymphocyte, e.g. a T cell, and shown that such substances may be used to produce APCs and lymphocytes, such as
50

55

5 regulatory T cells, capable of suppressing the activity of other APCs and lymphocytes, such as T cells.

10 Accordingly, the present invention also provides the use of a composition capable of
5 upregulating expression of an endogenous Notch or Notch ligand in an APC and/or lymphocyte in a method of producing APCs or regulatory lymphocytes, preferably T cells, capable of suppressing the activity of other APCs and/or lymphocytes. Typically, the
15 composition is used *in vitro/ex vivo* rather than *in vivo* and the resulting APCs/lymphocytes (T cells) subsequently administered to a patient.

10 Thus the present invention also provides a method of treating a patient suffering from a
20 disease characterised by inappropriate lymphocyte activity which method comprises administering to the patient a lymphocyte produced by the methods of the invention.

25 15 It is not necessary to incubate the APC and lymphocyte simultaneously: for example, the APC can be primed first in the presence of the antigen and substance capable of upregulating Notch or Notch ligand expression. When such a primed APC is contacted with a
30 lymphocyte, either *in vitro*, *ex vivo* or *in vivo*, tolerance to the antigen is induced in the lymphocyte.

20 Accordingly, the present invention further provides a method for producing an antigen
35 presenting cell (APC) capable of inducing in a lymphocyte tolerance to an allergen or antigen which method comprises contacting an APC with (i) a composition capable of upregulating expression of an endogenous Notch or Notch ligand in the APC or lymphocyte and (ii) the
40 25 allergen or antigen. Such APCs may be administered to a patient in a method of immunotherapy. Preferably the method is carried out *ex vivo* preferably using APCs or lymphocytes obtained from a human or animal patient suffering from an immune disorder or the recipient of a tissue graft/organ transfer.

45 30 Also provided is a method for producing *ex vivo* a lymphocyte having tolerance to an allergen or antigen which method comprises incubating an APC, produced as described above, with
50 the lymphocyte.

5 In one preferred embodiment step (i) comprises introducing a nucleic acid sequence into
the lymphocyte or APC, which is capable of upregulating expression of an endogenous
Notch or Notch ligand, preferably by expression of a polypeptide which is capable of
10 upregulating expression of an endogenous Notch or Notch ligand.

5

Preferably the nucleic acid sequence encodes a polypeptide selected from Noggin, Chordin,
Follistatin, Xnr3, fibroblast growth factors and derivatives, fragments, variants and
15 homologues thereof, and/or at least one immunosuppressive cytokine. Particularly preferred
examples of immunosuppressive cytokines for use in the present invention are IL-4, IL-10,
10 IL-13, TGF- β and FLT3 ligand.

20

Alternatively, the nucleic acid sequence is an antisense construct derived from a sense
nucleotide sequence encoding a polypeptide selected from a polypeptide capable of
downregulating Notch or Notch ligand expression, such as Toll-like receptors, a cytokine
25 such as IL-12, IFN- γ , TNF- α , or a growth factor such as a BMP or a BMP receptor and
15 activins.

30

In another preferred embodiment the composition is a chemical compound such as a
polypeptide which is exposed/incubated with the lymphocyte or APC. The agent should be
20 one which is capable of modulating Notch-Notch ligand interactions. In this embodiment the
polypeptide is preferably selected from from Noggin, Chordin, Follistatin, Xnr3, fibroblast
35 growth factors and derivatives, fragments, variants and homologues thereof, and/or at least
one immunosuppressive cytokine. Particularly preferred examples of immunosuppressive
cytokines for use in the present invention are IL-4, IL-10, IL-13, TGF- β and FLT3 ligand

40

25

Preferably when the composition is a receptor or a nucleic acid sequence encoding a receptor,
the receptor is activated. Thus, when the agent is a nucleic acid sequence, the receptor is
45 constitutively active when expressed.

45

30 As used herein, the terms protein and polypeptide may be assumed to be synonymous, protein
merely being used in a general sense to indicate a relatively longer amino acid sequence than
50 that present in a polypeptide.

55

5

The term "derivative" as used herein, in relation to proteins or polypeptides of the present invention includes any substitution of, variation of, modification of, replacement of, deletion of or addition of one (or more) amino acid residues from or to the sequence providing that the resultant protein or polypeptide possesses the capability of modulating Notch-Notch ligand interactions.

10

5

15

The term "variant" as used herein, in relation to proteins or polypeptides of the present invention includes any substitution of, variation of, modification of, replacement of, deletion of or addition of one (or more) amino acid residues from or to the sequence providing that the resultant protein or polypeptide possesses the capability of modulating Notch-Notch ligand interactions.

10

20

25

The term "analog" are used herein, in relation to the proteins or polypeptides of the present invention includes any peptidomimetic, that is, a chemical compound that possesses the capability of modulating Notch-Notch ligand interactions in a similar manner to the parent protein or polypeptide. These include compounds that may antagonise the expression or activity of a Notch-protein or Notch-ligand.

30

20 An agent may be considered to modulate Notch-Notch ligand interactions if it is capable of promoting the interaction of Notch with its ligands, preferably to an extent sufficient to provide therapeutic efficacy.

35

40

25 In a preferred embodiment the agent modulates Notch-Notch ligand interactions by being capable of upregulating expression of Notch or Notch ligand.

45

The expression "Notch-Notch ligand" as used herein means the interaction between a Notch family member and a ligand capable of binding to one or more such member. Thus by the expression "upregulating interaction of Notch or a Notch-ligand" we mean promoting the interaction of Notch in a lymphocyte or APC with a Notch ligand or promoting the interaction of a Notch ligand in a lymphocyte or APC with Notch. Preferably the lymphocyte is a T cell.

50

55

5 The term therapy are used herein should be taken to encompass diagnostic and prophylactic applications.

10 The present invention further provides a pharmaceutical composition comprising a primed APC and/or lymphocyte of the invention together with a pharmaceutically acceptable carrier or diluent.

15 Detailed Description of the Invention

20 Various preferred features and embodiments of the present invention will now be described by way of non-limiting example.

25 **A. Notch and Notch ligands**

30 An endogenous Notch ligand in the context of the present invention is a polypeptide encoded by the genome of a mammalian cell that is capable of being expressed by the mammalian cell. In particular the mammalian cell may be a haemopoietic cell such as a T cell or an antigen presenting cell. The endogenous Notch ligand is typically is capable of binding to a Notch receptor polypeptide present in the membrane of a variety of mammalian cell types, for example haemopoietic stem cells. At least four Notch receptors (Notch-1, Notch-2, Notch-3 and Notch-4) have been identified to date in human cells.

35 Particular examples of mammalian Notch ligands identified to date include the Delta family, for example Delta-1 (Genbank Accession No. AF003522 - *Homo sapiens*), Delta-3 (Genbank Accession No. AF084576 - *Rattus norvegicus*) and Delta-like 3 (*Mus musculus*), the Serrate family, for example Serrate-1 and Serrate-2 (WO97/01571, WO96/27610 and WO92/19734), Jagged-1 and Jagged-2 (Genbank Accession No. AF029778 - *Homo sapiens*), and LAG-2. Homology between family members is extensive. For example, human Jagged-2 has 40.6% identity and 58.7% similarity to Serrate.

40
45
50
55
30 Further homologues of known mammalian Notch ligands may be identified using standard techniques. By a "homologue" it is meant a gene product that exhibits sequence homology, either amino acid or nucleic acid sequence homology, to any one of the known Notch

5 ligands, for example as mentioned above. Typically, a homologue of a known Notch
ligand will be at least 20%, preferably at least 30%, identical at the amino acid level to the
corresponding known Notch ligand. Techniques and software for calculating sequence
10 homology between two or more amino acid or nucleic acid sequences are well known in
5 the art (see for example <http://www.ncbi.nlm.nih.gov> and Ausubel *et al.*, Current Protocols
in Molecular Biology (1995), John Wiley & Sons, Inc.)

15 Notch ligands identified to date have a diagnostic DSL domain (D. Delta, S. Serrate, L.
Lag2) comprising 20 to 22 amino acids at the amino terminus of the protein and between 3 to
10 8 EGF-like repeats on the extracellular surface. It is therefore preferred that homologues of
20 Notch ligands also comprise a DSL domain at the N-terminus and between 3 to 8 EGF-like
repeats on the extracellular surface.

25 In addition, suitable homologues will be capable of binding to a Notch receptor. Binding
15 may be assessed by a variety of techniques known in the art including *in vitro* binding
assays.

30 Homologues of Notch ligands can be identified in a number of ways, for example by
probing genomic or cDNA libraries with probes comprising all or part of a nucleic acid
20 encoding a Notch ligand under conditions of medium to high stringency (for example 0.03M
sodium chloride and 0.03M sodium citrate at from about 50°C to about 60°C). Alternatively,
35 homologues may also be obtained using degenerate PCR which will generally use primers
designed to target sequences within the variants and homologues encoding conserved amino
acid sequences. The primers will contain one or more degenerate positions and will be used
40 25 at stringency conditions lower than those used for cloning sequences with single sequence
primers against known sequences.

45 **B. Substances capable of upregulating endogenous Notch or Notch ligand
expression**

30 Substances that may be used to upregulate Notch ligand expression include polypeptides
50 that bind to and reduce or neutralise the activity of bone morphogenetic proteins (BMPs).

5 Binding of extracellular BMPs (Wilson and Hemmati-Brivanlou, 1997, Hemmati-Brivanlou
and Melton, 1997) to their receptors leads to down-regulated Delta transcription due to the
inhibition of the expression of transcription factors of the achaete/scute complex. This
10 complex is believed to be directly involved in the regulation of Delta expression. Thus, any
5 substance that inhibits BMP expression and/or inhibits the binding of BMPs to their receptors
may be capable of producing an increase in the expression of Notch ligands such as Delta
and/or Serrate. Particular examples of such inhibitors include Noggin (Valenzuela *et al.*,
15 1995), Chordin (Sasai *et al.*, 1994), Follistatin (Iemura *et al.*, 1998), Xnr3, and derivatives
and variants thereof. Noggin and Chordin bind to BMPs thereby preventing activation of their
10 signalling cascade which leads to decreased Delta transcription. Consequently, increasing
Noggin and Chordin levels may lead to increase Notch ligand, in particular Delta, expression.

20
Furthermore, any substance that upregulates expression of transcription factors of the
achaete/scute complex may also upregulate Notch ligand expression.

25 15
Other suitable substances that may be used to upregulate Notch ligand expression include
transforming growth factors such as members of the fibroblast growth factor (FGF) family.
The FGF may be a mammalian basic FGF, acidic FGF or another member of the FGF
30 family such as an FGF-1, FGF-2, FGF-3, FGF-4, FGF-5, FGF-6, FGF-7. Preferably the
20 FGF is not acidic FGF (FGF-1; Zhao *et al.*, 1995). Most preferably, the FGF is a member
of the FGF family which acts by stimulating the upregulation of expression of a Serrate
polypeptide on APCs. The inventors have shown that members of the FGF family can
35 upregulate Serrate-1 gene expression in APCs.

40 25
Immunosuppressive cytokines may also be used to upregulate Notch ligand expression.
Examples include members of the TGF- β family such as TGF- β -1 and TGF- β -2, and
interleukins such as IL-4, IL-10 and IL-13, and FLT3 ligand.

45 30
The inventors have shown that: members of the TGF- β family can upregulate Notch,
particularly Notch 1, expression in CD4+ T cells;

50
IL-10 can upregulate serrate, particularly Serrate 1, gene expression in dendritic cells;

5

IL-10 can upregulate Notch, Delta and Serrate, particularly Notch 2, Notch 4, Delta 1 and Serrate 1, gene expression in naive B cells; and

10

5 IL-10 can upregulate Serrate, particularly Serrate 1, gene expression in mature DCs.

15

Generally the inventors have shown that selected cytokines affect different receptors/ligands in different ways, and that this also varies between cell type and tissue.

20

10 The substance capable of upregulating expression of Notch or a Notch ligand may be selected from polypeptides and fragments thereof, linear peptides, cyclic peptides, synthetic and natural compounds including low molecular weight organic or inorganic compounds. The substances capable of upregulating expression of a Notch ligand may be derived from a biological material such as a component of extracellular matrix. Suitable extracellular
25 15 matrix components are derived from immunologically privileged sites such as the eye. For example aqueous humour or components thereof may be used.

30

Polypeptide substances such as Noggin, FGFs and TGF- β may be purified from mammalian cells, obtained by recombinant expression in suitable host cells or obtained
20 commercially. Alternatively, nucleic acid constructs encoding the polypeptides may be introduced into APCs and/or lymphocytes (T cells) by transfection using standard techniques or viral infection/transduction. As a further example, overexpression of Notch
35 or Notch ligand, such as Delta or Serrate, may be brought about by introduction of a nucleic acid construct capable of activating the endogenous gene, such as the Serrate or
40 25 Delta gene. In particular, gene activation can be achieved by the use of homologous recombination to insert a heterologous promoter in place of the natural promoter, such as the Serrate or Delta promoter, in the genome of the APC or lymphocyte (T cell).

45

It is particularly preferred to use combinations of substances, for example a combination of
30 at least two substances. In a preferred embodiment, an immunosuppressive cytokine is used in combination with another substance capable of upregulating Notch ligand expression. Other examples of preferred combinations include at least one substance capable of
50

55

5 upregulating Serrate expression (such as FGF), preferably in an APC, together with at least one substance capable of upregulating Delta expression (such as Noggin or Chordin), preferably in a T cell. Alternatively, a preferred combination comprises at least one
10 substance which acts via inhibition of binding of BMPs to their receptors together with at least one substance which has a different mode of action.

15 Preferably, the composition, preferably a nucleic acid sequence, for use in the present invention is capable of upregulating Serrate and Delta, preferably Serrate 1 and Serrate 2 as well as Delta 1 and Delta 3 expression in APCs such as dendritic cells.

20 Preferably, the substance for use in the present invention is capable of upregulating Serrate expression in APCs such as dendritic cells. In particular, the substance may be capable of upregulating Serrate expression but not Delta expression in APCs. Alternatively, the substance for use in the present invention is capable of upregulating Delta expression in T
25 15 cells such as CD4⁺ helper T cells or other cells of the immune system that express Delta (for example in response to stimulation of cell surface receptors). In particular, the substance may be capable of upregulating Delta expression but not Serrate expression in T cells. In a particularly preferred embodiment, the substance is capable of upregulating
30 Notch ligand expression in both T cells and APC, for example Serrate expression in APCs and Delta expression in T cells.

35 Suitable substances for use according to the present invention may be conveniently identified using a simple screening procedure. In one such assay procedure, lymphocytes, such as T cells, or APCs in culture may be contacted with a candidate substance and the
40 25 effect on expression of an endogenous Notch ligand, such as Delta or Serrate, determined, for example by (i) measuring transcription initiated from the gene encoding the Notch ligand as described in the Examples or by quantitative-reverse transcriptase-polymerase chain reaction (RT-PCR); (ii) detecting Notch ligand protein by techniques such as
45 Western blotting of cell extracts, immunohistochemistry or flow cytometry; and/or (iii) 30 functional assays such as cell adhesion assays.

50 The present invention also relates to modification of Notch-protein expression or presentation on the cell membrane or signalling pathways. Agents that enhance the

5 presentation of a fully functional Notch-protein on the lymphocyte or APC surface include
matrix metalloproteinases such as the product of the Kuzbanian gene of *Drosophila* (Dkuz
et al (1997)) and other ADAMALYSIN gene family members.

10 5 In more detail, whether a substance can be used for modulating Notch-Notch ligand
expression may be determined using suitable screening assays.

15 Screening assays for the detection of increased Notch, Notch ligand expression and/or
processing include:

10

20 Notch-Notch ligand expression may be assessed following exposure of isolated cells to test
compounds in culture using for example:

25 15 (a) at the protein level by specific antibody staining using immunohistochemistry or flow
cytometry.

30 (b) at the RNA level by quantitative - reverse transcriptase-polymerase chain reaction (RT-
PCR). RT-PCR may be performed using a control plasmid with in-built standards for
20 measuring endogenous gene expression with primers specific for Notch 1 and Notch 2,
Serrate 1 and Serrate 2, Delta 1 and Delta 2 and Delta 3. This construct may be modified as
new ligand members are identified.

35 (c) at the functional level in cell adhesion assays.

40 25 Increased Notch ligand or Notch expression should lead to increased adhesion between
cells expressing Notch and its ligands. Test cells will be exposed to a particular treatment
in culture and radiolabelled or flourescein labelled target cells (transfected with
Notch/Notch ligand protein) will be overlaid. Cell mixtures will be incubated at 37°C for
45 2 hours. Nonadherent cells will be washed away and the level of adherence measured by
30 the level of radioactivity/immunofluorescence at the plate surface.

50 Using such methods it is possible to detect compounds or Notch-ligands that affect the
expression or processing of a Notch-protein or Notch-ligand. The invention also relates to

5 compounds, or Notch-ligands detectable by these assays methods, and also to their use in
the methods of the present invention.

10 These procedures may also be used to identify particularly effective combinations of
5 substances for use according to the present invention.

15 **C. Antigen Presenting Cells and Lymphocytes**

15 Antigen-presenting cells (APCs) for use in the present invention may be "professional"
10 antigen presenting cells or may be another cell that may be induced to present antigen to T
cells. Alternatively a APC precursor may be used which differentiates or is activated under
20 the conditions of culture to produce an APC. The APC may be isolated from a patient, or
recipient of the immunotherapy or from a donor individual or another individual.
Preferably the APC or precursor is of human origin. If the APC or precursor APC is from a
25 15 different individual to the T cells, the donor APC may also serve as the source of antigen.

30 APCs include dendritic cells (DCs) such as interdigitating DCs or follicular DCs,
Langerhans cells, PBMCs, macrophages, B-lymphocytes, T-lymphocytes, or other cell
types such as epithelial cells, fibroblasts or endothelial cells, activated to express a MHC
20 molecule (Class I or II) on their surfaces. Precursors of APCs include CD34⁺ cells,
monocytes, fibroblasts and endothelial cells. The APCs or precursors may be modified by
35 the culture conditions or may be genetically modified, for instance by transfection of one or
more genes encoding proteins which play a role in antigen presentation. Such proteins
include MHC molecules (Class I or Class II), CD80, CD86, or CD40. Most preferably
25 DCs or DC-precursors are included as a source of APCs.

40 The APC or precursor APC may be provided by a cell proliferating in culture such as an
established cell line or a primary cell culture. Examples include hybridoma cell lines,
45 L-cells and human fibroblasts such as MRC-5. Cell lines may conveniently be used in the
30 screening procedures described above.

50 Dendritic cells (DCs) can be isolated/prepared by a number of means, for example they can
either be purified directly from peripheral blood, or generated from CD34⁺ precursor cells

5 for example after mobilisation into peripheral blood by treatment with GM-CSF, or directly
from bone marrow. From peripheral blood, adherent precursors can be treated with a
GM-CSF/IL-4 mixture (Inaba *et al.*, 1992), or from bone marrow, non-adherent CD34⁺
10 cells can be treated with GM-CSF and TNF- α (Caux *et al.*, 1992). DCs can also be
5 routinely prepared from the peripheral blood of human volunteers, similarly to the method
of Sallusto and Lanzavecchia (1994) using purified peripheral blood mononucleocytes
(PBMCs) and treating 2 hour adherent cells with GM-CSF and IL-4. If required, these may
15 be depleted of CD19⁺ B cells and CD3⁺, CD2⁺ T cells using magnetic beads (see Coffin *et al.*, 1998). Culture conditions may include other cytokines such as GM-CSF or IL-4 for the
10 maintenance and, or activity of the dendritic cells or other antigen presenting cells.

20 Where lymphocytes are to be used they are preferably T cells or B cells. T cells are most preferred.

25 15 Where T cells or B cells are to be used in *ex vivo* methods of inducing immunotolerance, the T cells or B cells for use in the invention are typically isolated from an individual suffering from a disease of the immune system or a recipient for a transplant operation or from a related or unrelated donor individual. T cells or B cells may be obtained from blood
30 or another source (such as lymph nodes, spleen, or bone marrow) and may be enriched or
20 purified by standard procedures. The T cells or B cells may be used in combination with other immune cells, obtained from the same or a different individual. Alternatively whole
35 blood may be used or leukocyte enriched blood or purified white blood cells as a source of T cells, B cells and other cell types. It is particularly preferred to use helper T cells (CD4⁺). Alternatively other T cells such as CD8⁺ cells may be used.

40 25 Where T cells or B cells are to be used in *in vitro* screening procedures, it may be convenient to use cell lines, such as T cell hybridomas.

45 30 Thus, it will be understood that the term "antigen presenting cell or the like" as used herein is not intended to be limited to APCs. The skilled man will understand that any vehicle capable of presenting to the T cell population may be used, for the sake of convenience the term APCs is used to refer to all these. As indicated above, preferred examples of suitable
50

5 APCs include dendritic cells, L cells, hybridomas, fibroblasts, lymphomas, macrophages, B
cells or synthetic APCs such as lipid membranes.

10 **E. Antigen and Allergens**

5 An antigen may be any substance that can be recognised generally as foreign, by the
immune system, and is generally recognised by an antigen receptor. Preferably the antigen
15 of the present invention is an immunogen. An allergic response occurs when the host is re-
exposed to an antigen that it has encountered previously.

10 The immune response to antigen is generally either cell mediated (T cell mediated killing)
20 or humoral (antibody production via recognition of whole antigen). The pattern of
cytokine production by TH cells involved in an immune response can influence which of
these response types predominates: cell mediated immunity (TH1) is characterised by high
25 15 IL-2 and IFN γ but low IL-4 production, whereas in humoral immunity (TH2) the pattern is
low IL-2 and IFN γ but high IL-4, IL-5, IL-10. Since the secretory pattern is modulated at
the level of the secondary lymphoid organ or cells, then pharmacological manipulation of
30 the specific TH cytokine pattern can influence the type and extent of the immune response
generated.

20 The TH1-TH2 balance refers to the interconversion of the two different forms of helper T
35 cells. The two forms have large scale and opposing effects on the immune system. If an
immune response favours TH1 cells, then these cells will drive a cellular response, whereas
TH2 cells will drive an antibody-dominated response. The type of antibodies responsible
40 25 for some allergic reactions is induced by TH2 cells.

The present invention has uses in relation to both responses.

45 The antigen or allergen may be a peptide, polypeptide, carbohydrate, protein, glycoprotein,
30 or more complex material containing multiple antigenic epitopes such as a protein
complex, cell-membrane preparation, whole cells (viable or non-viable cells), bacterial
50 cells or virus/viral component. In particular, it is preferred to use antigens known to be

5 associated with auto-immune diseases such as myelin basic protein (associated with
multiple sclerosis), collagen (associated with rheumatoid arthritis), and insulin (diabetes),
or antigens associated with rejection of non-self tissue such as MHC antigens. Where
10 primed APCs/ T cells of the present invention are to be used in tissue transplantation
5 procedures, antigens will be obtained from the tissue donor.

The antigen or allergen moiety may be, for example, a synthetic MHC-peptide complex i.e. a
15 fragment of the MHC molecule bearing the antigen groove bearing an element of the antigen.
Such complexes have been described in Altman *et al.*, 1996.

10

20 **F. Preparation of Primed APCs and Lymphocytes**

1. Preparation of Primed APCs *ex vivo* in the absence of lymphocytes

25 15 APCs as described above are cultured in a suitable culture medium such as DMEM or other
defined media, optionally in the presence of fetal calf serum. Cytokines, if present, are
typically added at up to 1000 U/ml. Optimum concentrations may be determined by
30 titration. One or more substances capable of upregulating Notch or Notch ligand expression
are then typically added to the culture medium together with the antigen of interest. The
20 antigen may be added before, after or at substantially the same time as the substance(s). Cells
are typically incubated with the substance(s) and antigen for at least one hour, preferably at
35 least 3 hours, at 37°C. If required, a small aliquot of cells may be tested for upregulation of
Notch or Notch ligand expression as described above. Alternatively, cell activity may be
measured by the inhibition of T cell proliferation as described in WO98/20142. APCs
40 25 transfected with a nucleic acid construct directing the expression of, for example Serrate, may
be used as a control.

As discussed above, polypeptide substances may be administered to APCs by introducing
45 nucleic acid constructs/viral vectors encoding the polypeptide into cells under conditions that
30 allow for expression of the polypeptide in the APC. Similarly, nucleic acid constructs
encoding antigens may be introduced into the APCs by transfection, viral infection or viral
50 transduction.

5 The resulting APCs that express increased levels of a Notch ligand and are presenting antigen on their cell surface complexed with MHC are now ready for use. For example, they may be prepared for administration to a patient or incubated with T cells *in vitro* (*ex vivo*) to induce immunotolerance in the T cells as described in WO98/20142.

10 5

2. Preparation of Regulatory T cells (and B cells) *ex vivo*

15 The techniques described below are described in relation to T cells, but are equally applicable to B cells. The techniques employed are essentially identical to that described for APCs alone except that T cells are generally co-cultured with the APCs. However, it may be preferred to prepare primed APCs first and then incubate them with T cells. For example, once the primed APCs have been prepared, they may be pelleted and washed with PBS before being resuspended in fresh culture medium. This has the advantage that if, for example, it is desired to treat the T cells with a different substance(s) capable of upregulating Notch or Notch ligand expression and/or cytokine to that used with the APC, then the T cell will not be brought into contact with the different substance(s) used to upregulate Notch or Notch ligand expression in the APC. Alternatively, the T cell may be incubated with the substance(s)/cytokine first to induce Notch or Notch ligand expression, washed, resuspended and then incubated with the primed APC in the absence of both the substance(s) used to upregulate APC Notch ligand expression and the substance(s) used to upregulate Notch or Notch ligand expression in the T cell. Once primed APCs have been prepared, it is not always necessary to administer any substances to the T cell since the primed APC is itself capable of inducing immunotolerance leading to increased Notch or Notch ligand expression in the T cell, presumably via Notch/Notch ligand interactions between the primed APC and T cell.

25 40

Incubations will typically be for at least 1 hour, preferably at least 3 or 6 hours, in suitable culture medium at 37°C. The progress of induction of Notch or Notch ligand expression may be determined for a small aliquot of cells using the methods described above. T cells transfected with a nucleic acid construct directing the expression of, for example Delta, may be used as a control. Induction of immunotolerance may be determined by subsequently challenging T cells with antigen and measuring IL-2 production compared with control cells not exposed to APCs.

45 50

55

5 Primed T cells or B cells may also be used to induce immunotolerance in other T cells or B
cells in the absence of APCs using similar culture techniques and incubation times.
Generally, the addition of substances capable of upregulating Notch or Notch ligand
10 expression is not required at this stage but they may be added if desired, together with
5 immunosuppressive cytokines.

G. Transgenic animals

15 The present invention also relates to cell lines or transgenic animals which are capable of
10 expressing or overexpressing Notch, a Notch ligand or at least one agent useful in the
present invention. Preferably the cell line or animal expresses or overexpresses Notch,
20 Delta or Serrate.

25 The present invention additionally relates to cell lines or transgenic animals which are
15 capable of expressing or overexpressing at least one polypeptide which is capable of
promoting Notch-Notch ligand interactions. Such agents have been described above and for
the avoidance of doubt are specifically incorporated herein by reference.

30 The present invention further relates to cell lines or transgenic animals which are capable
20 of expressing or overexpressing at least one polypeptide which is capable of enhancing
Notch-Notch ligand interactions. Agents that enhance the presentation of a fully functional
35 Notch-protein on the lymphocyte or APC surface include matrix metalloproteinases such as
the product of the Kuzbanian gene of *Drosophila* (Dkuz *et al.*, (1997) and other
ADAMALYSIN gene family members. Suitable agents that influence expression of
25 Notch-ligands include agents that affect the expression of Delta and/or Serrate genes. For
instance, for Delta expression, any agent that inhibits the binding of BMPs to their
40 receptors is capable of producing an increase in the expression of Delta and/or Serrate.
Such agents include Noggin, Chordin, Follistatin, FGFs, Fringe and derivatives and
45 variants thereof.

30 The transgenic animal is typically a vertebrate, more preferably a rodent, such as a rat or a
50 mouse, but also includes other mammals such as human, goat, pig or cow etc.

5 Such transgenic animals are useful as animal models of disease and in screening assays for
new useful compounds. By specifically expressing one or more polypeptides, as defined
above, the effect of such polypeptides on the development of disease can be studied.
10 Furthermore, therapies including gene therapy and various drugs can be tested on
5 transgenic animals. Methods for the production of transgenic animals are known in the
art. For example, there are several possible routes for the introduction of genes into
embryos. These include (i) direct transfection or retroviral infection of embryonic stem
15 cells followed by introduction of these cells into an embryo at the blastocyst stage of
development; (ii) retroviral infection of early embryos; and (iii) direct microinjection of
10 DNA into zygotes or early embryo cells.

20 The present invention also includes stable cell lines for use as disease models for testing or
treatment. A stable cell line will contain a recombinant gene or genes, also known herein
as a transgene.

25 15 A cell line containing a transgene, as described herein, is made by introducing the
transgene into a selected cell line according to one of several procedures known in the art
for introducing a foreign gene into a cell.

30 20 The sequences encoding the inhibitors and enhancers of Notch-Notch ligand interactions as
well as Notch or a Notch ligand itself are operably linked to control sequences, including
35 promoters/enhancers and other expression regulation signals.

40 25 The promoter is typically selected from promoters which are functional in mammalian
cells, although prokaryotic promoters and promoters functional in other eukaryotic cells
may be used. The promoter is typically derived from promoter sequences of viral or
eukaryotic genes. For example, it may be a promoter derived from the genome of a cell in
45 which expression is to occur. With respect to eukaryotic promoters, they may be promoters
that function in a ubiquitous manner (such as promoters of α -actin, β -actin, tubulin) or,
30 alternatively, a tissue-specific manner (such as promoters of the genes for pyruvate kinase).
Tissue-specific promoters specific for lymphocytes, dendritic cells, skin, brain cells and
epithelial cells within the eye are particularly preferred, for example the CD2, CD11c,
50 keratin 14, Wnt-1 and Rhodopsin promoters respectively. Preferably the lung epithelial

5 cell promoter SPC is used. They may also be promoters that respond to specific stimuli,
for example promoters that bind steroid hormone receptors. Viral promoters may also be
used, for example the Moloney murine leukaemia virus long terminal repeat (MMLV LTR)
10 promoter, the rous sarcoma virus (RSV) LTR promoter or the human cytomegalovirus
5 (CMV) IE promoter.

It may also be advantageous for the promoters to be inducible so that the levels of
15 expression of the heterologous gene can be regulated during the life-time of the cell.
Inducible means that the levels of expression obtained using the promoter can be regulated.

10 In addition, any of these promoters may be modified by the addition of further regulatory
sequences, for example enhancer sequences. Chimeric promoters may also be used
20 comprising sequence elements from two or more different promoters described above.

25 15 **H. Therapeutic Uses**

We have shown that APCs and lymphocytes expressing Notch and Notch ligands are
30 capable of efficiently transferring infectious tolerance to the chosen antigen or antigens
when transferred into the patient for the treatment of a disease characterised by
20 inappropriate lymphocyte activity, such as Th1 or Th2 cell activity. The APCs and/or
lymphocytes may thus be used to treat an ongoing immune response (such as an allergic
condition or an autoimmune disease) or may be used to generate tolerance in an
35 immunologically lymphocytes cells of the present invention may be used in therapeutic
methods for both treating and preventing diseases characterised by inappropriate lymphocyte
25 activity in animals and humans. The APCs and/or lymphocytes may be used to confer
tolerance to a single antigen or to multiple antigens.

Typically, APCs and/or lymphocytes are obtained from the patient or donor and primed as
45 described above before being returned to the patient (*ex vivo* therapy).

30 Particular conditions that may be treated or prevented include multiple sclerosis,
rheumatoid arthritis, diabetes, allergies, asthma, and graft rejection. The present invention
50 may also be used in organ transplantation or bone marrow transplantation.

5

I. Administration

10

5 Primed APCs/lymphocytes of the present invention for use in immunotherapy are typically formulated for administration to patients with a pharmaceutically acceptable carrier or diluent to produce a pharmaceutical composition. Suitable carriers and diluents include isotonic saline solutions, for example phosphate-buffered saline. The composition may be formulated for parenteral, intramuscular, intravenous, intra-peritoneal, injection, intranasal inhalation, lung inhalation, intradermal, intra-articular, intrathecal, or via the alimentary tract (for example, via the Peyers patches).

15

20

Cells and pharmaceutical comprising cells of the invention are typically administered to the patient by intramuscular, intraperitoneal or intravenous injection, or by direct injection into the lymph nodes of the patient, preferably by direct injection into the lymph nodes.

25

15 Typically from 10^4 to 10^8 treated cells, preferably from 10^5 to 10^7 cells, more preferably about 10^6 cells are administered to the patient.

30

The routes of administration and dosages described are intended only as a guide since a skilled practitioner will be able to determine readily the optimum route of administration and dosage for any particular patient depending on, for example, the age, weight and condition of the patient.

35

The present invention will now be described by way of examples which are intended to be illustrative only and non-limiting, and by reference to the accompanying Figures in which

40

25 Figures 1, 2, 3 and 4 show the results of Examples 8, 9, 10 and 11 respectively.

EXAMPLES

45

Materials and Methods

30

Construction of Retroviruses expressing Delta or Serrate

50

55

5 A cDNA encoding human Serrate-1 is inserted as a *Bam*HI – *Eco*RI fragment into the
retroviral vector pBABEneo (Morgenstern and Land, 1990) using standard techniques,
such that the cDNA is expressed from the retroviral promoter element. Transducing vector
10 particles can be produced by transfection of this construct into a suitable amphotropic
5 packaging cell line such as PA317 (ATCC catalogue number CRL-9078), FLYA13 or
FLYRD114 (Cosset *et al.*, 1995). Alternatively a permanent cell line containing the vector
genome and gag-pol gene is constructed by transfection of Serrate-pBABE vector into
15 PA327 cells. Production of replication – deficient pseudotyped vector particles is initiated
by transient transfection with a plasmid encoding the G-protein from Vesicular Stomatitis
10 Virus (VSV) under the control of the hCMV major immediate early promoter-enhancer.

20 A retroviral vector expressing mouse Delta-1 is prepared in a similar fashion.

Preparation of Dendritic Cells from Mouse Spleen

25 15 Single cell suspensions are prepared from the spleens of mice. Rapidly adhering cells are
isolated by culture for 2-3 h at 37°C in plastic tissue culture flasks. Non-adherent cells are
removed by extensive washing and 50 ng/ml mouse GM-CSF in culture medium is added
for 24 h. Culture medium is RPMI 1640 with 2% FCS, 50 mM 2-mercaptoethanol, and
30 optionally 0.02 mg/ml Penicillin and Streptomycin.

20

Preparation of T cells from Mouse

35 T cells are purified from blood or lymph nodes by positive selection on magnetic antibody-
coated beads specific for particular cell types (MACS columns) using methods provided by
the manufacturer (Miltenyi Biotech) as follows.

25

40 Lymph nodes are removed and single cell suspensions prepared in tissue culture medium
(10^8 cells in 0.4 ml RPMI 1640 with 10% FCS). Cells are incubated at 4°C for 15 min,
passed over the MACS separation column, washed and collected. CD4-positive cells are
45 enriched by negative selection on anti-CD8 antibody-coated magnetic beads.

30

Determination of Serrate and Delta expression.

50 After various incubation times, cells (1.5×10^6) are harvested, pelleted and frozen. RNA is
prepared from cell pellets by homogenisation in guanidium thiocyanate solution followed by

5 CsCl density centrifugation. 1 µg RNA is converted into cDNA using an oligo dT primer. Of
the resultant cDNA, 1/20th was used in PCR (40 cycles) using primers specific for the human
delta homologue or the human serrate homologue.

10 5 RT-PCR is performed using an Access RT-PCR kit (Promega). 50 ng RNA is used in each
reaction together with Serrate-1 gene specific oligonucleotide primers (50 pmol) under
conditions according to the manufacturer's instructions (T_m for the Serrate oligonucleotides
15 is 58°C).

10 The sequence of the "forward" Serrate-1 primer is:
20 5'-GGCTGGGAAGGAACAACCTG-3'

The Serrate- "reverse" primer is:
25 5'-GGTAGCCATTGATCTCATCCAC-3'

15 Primers specific for Delta are:
25 5'-GATTCTCCTGATGACCTCGC-3'
30 5'-GTGTTCGTCACACACGAAGC-3'

20 PCR samples were analysed by gel electrophoresis.

35 *Determination of Notch, Serrate and Delta Expression*

In the Examples 8-11 the following primers were used as indicated:

Murine Notch1 Accession No. Z11886
40 25 Forward primer (FP) 5'-TGTTAATGAGTGCATCTCCAACCC-3'
Reverse primer (RP) 5'-CATTCGTAGCCATCAATCTTGTC-3'

Murine Notch2 Accession No. D32210
45 FP 5'-CAGAGGAATAGCAAGACGTGCAAG-3'
30 RP 5'-GATGAAGAACAGGATGATGACAACAG-3'

50 Murine Notch4 Accession No. U43691

5 FP 5'-CTACTGCCACAAGTAGCTGG-3'
 RP 5'-CTCGGAGATAGCGTGACTGG-3'

Murine Serrate1 (Jagged 1) Shimizu *et al* 1999

10 5 FP 5'-GGGGTCACTGTCAGAATGA-3'
 RP 5'-AGATA'ACCGCACCCCTTCAG-3'

Murine Serrate2 (Jagged 2)

15 FP 5'-ATCTGCGAGGACCTGGTGGAT-3'
10 RP 5'-TATACCAGAGGGTGGCACA-3'

Murine Deltal Accession No. X80903

20 FP 5'-GACTCTCCCGATGACCTC-3'
 RP 5'-GATGCACTCATCGCAGTAG-3'

25 15

Example 1 - Assays to determine to identify substances that upregulate Notch ligand expression.

30

20 Dendritic cells (DCs) are the primary antigen presenting cell in the immune system and are critical for stimulating T cell responses. DCs are obtained from the spleen of mice and transferred to flasks containing tissue culture medium (RPMI 1640 with 10% fetal calf serum added). Cytokines (eg IL-4 and GM-CSF) are added as appropriate.

35

40

25 Cells are then transferred into 12-well tissue culture trays. To each well is added a different candidate upregulator of Notch ligand expression. Delta and Serrate expression is monitored at various time points by removing an aliquot of cells and determining induction of Delta and Serrate expression by PCR.

45

30 Similar procedures are also carried out using a T cell hybridoma cell line and T cells obtained from mice as described in the materials and methods section.

50

55

5 Example 2 - Preparation of Primed Dendritic Cells

10 DCs are obtained from the spleen of mice as in Example 1 and divided into two cultures. The first culture is transfected with a retrovirus allowing expression of the full length Serrate-1 protein to serve as a positive control. The first culture is then pulsed with the HDM peptide p110-131 for 3 hours at 37°C. The second culture is split up into several tissue culture plate wells and to each well is added a different upregulator of Notch ligand expression identified in Example 1. These wells are then also pulsed with the HDM peptide p110-131 for 3 hours at 37°C

15
20 The DCs are then washed and used to immunise naive mice subcutaneously using 10^5 cells/mouse. After 7 days the draining LNCs are recovered and restimulated in culture with peptide at 4×10^5 cells/well. Since the mice were only immunised with peptide-pulsed DCs this gives us a measure of the ability of these cells to prime an immune response.

25 Example 3 - Upregulation of Serrate expression in antigen presenting cells prevents T cell responses.

30 An influenza-reactive human T cell clone HA1.7 is mixed with peptide HA306-318 (1.0 µg/ml) in the presence of L cells expressing HLA-DRB1*0101 (as antigen presenting cells), using 2×10^4 of each cell type. The L cells have been preincubated with one or more substances identified as being capable of upregulating Serrate expression in APCs for 6 hours. The proliferative response is measured after 72 hours following addition of tritiated thymidine for the last 8 hours of culture.

40 Example 4 - Serrate upregulated antigen presenting cells induce regulatory T cells that can block the response of normal T cells.

45 An influenza-reactive human T cell clone HA1.7 is mixed with peptide HA306-318 and L cells (expressing DRB1*0101 as antigen presenting cells) in the presence of 2% IL-2. The L cells have been preincubated with one or more substances identified as being capable of upregulating Serrate expression in APCs for 6 hours. After 7 days in culture, the HA1.7 cells

5 were harvested, washed and irradiated before being mixed with fresh HA1.7 (using 2×10^4
each population). Cells are cultured for a further 2 days before being stimulated with peptide
(1.0 $\mu\text{g/ml}$) + normal APCs (DRB1*0101 PBMCs). The proliferative response is measured
10 after 72 hours following addition of tritiated thymidine for the last 8 hours of culture.

5 The results show the ability of cells tolerised by upregulation of Serrate to pass on their
tolerance to a naive cell population (infectious/bystander tolerance).

15 **Example 5 - Preparation of regulatory T cells *ex vivo* using primed APCs**

10 Primed dendritic cells are produced using the same method as in Example 2. These cells
are then washed, pelleted and resuspended in fresh culture medium. T cells obtained from
the mouse host are then incubated with the primed dendritic cells for up to 6 hours.
Aliquots of cells are taken at regular intervals and Delta and Serrate expression measured.

25 15 Helper T cells are separated from the other cells using magnetic beads specific for CD4
prior to pelleting and RNA extraction.

30 Induction of immunotolerance in the T cells is also measured in a functional assay. HDM
peptide p110-131 added to the cell culture and the cells cultured for 24 hours. Supernatant
20 fluids are then collected and assayed for IL-2 (a major T cell growth factor) content.

35 **Example 6 - Delta-upregulated primed T cells are able to inhibit the development of
immunity to peptide 110-131 antigen in animals.**

40 25 1×10^7 primed regulatory T cells generated by the method described in Example 4 are injected
into C57 BL mice i.p. The mice are also immunised with 50 μg Der p1 emulsified in
Complete Freund's Adjuvant (CFA) sub-cutaneously. After 7 days the draining lymph node
cells are collected and cultured at 4×10^5 cells/well with Der p1 (10 $\mu\text{g/ml}$) or peptide 110-
45 131 of Der p1 (10 $\mu\text{g/ml}$). Cultures are incubated at 37°C for 72 hours and tritiated
30 thymidine added for the final 8 hours of culture.

50

55

5 The results show that the primed regulatory T cells inhibit the development of an immune response to the Der p1 antigen in the immunised mouse.

Example 7 – Treatment of patients undergoing Bone Marrow Transplantation

10 5 Donor individuals for the bone marrow transplantation procedure are selected from an appropriate category (live related; MHC-matched un-related or unmatched); DCs are isolated from the donor by a suitable method (eg as described in US-A-5789148) between 15 14 days prior to, and 3 days after, transplantation. DCs are maintained in culture in tissue culture medium eg RPMI-1640 supplemented with up to 10% autologous or ABO human serum). Inducers of Notch-ligand expression are added for the appropriate time (between 20 3 h and 2 days). Cytokines are also added as required (eg IL-4 and GM-CSF).

DCs may be similarly prepared from the transplant recipient if required.

25 15 Lymphocytes are obtained by an appropriate method (e.g. according to the procedures described in US-A-4663058) from the donor and/or recipient. T cells may be enriched by standard methods including antibody-mediated separation. Cells are cultured in RPMI- 30 1640 with serum (autologous or ABO human serum) together with DCs.

20 T cells and DCs are then transferred to the transplant recipient by infusion at a suitable time, between 14 days before and 3 days after transplantation.

35 Other modifications of the present invention will be apparent to those skilled in the present art.

40 **Example 8 - Modulation of the expression of Notch receptors and ligands on activated murine CD4+ T cells in response to inflammatory and immunosuppressive stimuli**

45 30 CD4+ T cells were separated by MACS column from the spleens of naive BALB/C mice. Cells (2×10^6 /ml) were cultured for 48 hours in tissue culture medium (RPMI 1640) supplemented with penicillin/streptomycin, L-glutamine and 5% foetal calf serum) at 37°C and activated by 5µg/ml plated anti-CD3 and 5µg/ml soluble anti-CD28 antibodies alone or

5 together with interleukin 10, IL-10 (80ng/ml), lipopolysaccharide, LPS (10µg/ml),
 transforming growth factor β , TGF- β (10ng/ml) or interferon γ , IFN- γ (10ng/ml). Cells
 were collected and centrifuged at 1500 rpm and mRNA was isolated using Oligotex
 10 following the manufacturer's instructions, transcribed into cDNA and analysed by real time
 5 PCR (TaqMan, ABI) following the manufacturer's instructions.

Results The results are illustrated in Figure 1 reveal the following changes in
 15 transcript levels for activated CD4+ T cells:

	Serrate1	IL-10 no change	Delta1	IL-10 no change
10		TGF- β decreased		TGF- β decreased
20		LPS decreased		LPS decreased
		IFN- γ no change		IFN- γ no change
25	Notch1	IL-10 decreased		
15		TGF- β decreased		
		LPS decreased		
		IFN- γ no change		

30
 20 **Example 9 - Modulation of the expression of Notch receptors and ligands on naive
 murine dendritic cells (DCs) in response to inflammatory and immunosuppressive
 35 stimuli**

CD11c+ DCs were MACS separated from the spleens of naive BALB/C mice and
 incubated at 37°C for 24 hours in medium alone or together with IL-10 (50 and 100ng/ml),
 40 25 lipopolysaccharide (LPS; 1, 5 and 10µg/ml) or TGF- β (1 and 10ng/ml). Cells were
 collected and centrifuged at 1500 rpm and mRNA was isolated using Oligotex following
 the manufacturer's instructions. RT-PCR was performed using an Access RT-PCR kit
 (Promega). One µg of total RNA was used in each reaction together with specific
 45 oligonucleotide primers (50 pmol) for the following genes as indicated under conditions
 30 according to the manufacturer's instructions. PCR was performed using a Hybaid machine,
 dynazyme II polymerase, 1.5 mM Mg, 28-35 cycles at an annealing temperature between
 50 56-63°C.

5

Results: The results are illustrated in Figure 2 and reveal the following changes in transcript levels for naive splenic CD11c+ DCs

10

Serrate1	IL-10 increased	Notch2	IL-10 no change
	TGF- β no change		TGF- β no change
	LPS decreased		LPS increased

5

15

Example 10 - Modulation of the expression of Notch receptors and ligands on naive B cells in response to inflammatory and immunosuppressive stimuli

10

20

B cells were MACS separated from the spleens of naive BALB/C mice and incubated at 37°C for 24 hours in medium alone or together with IL-10 (80ng/ml) or lipopolysaccharide (LPS; 5 and 10 μ g/ml). Cells were collected and centrifuged at 1500 rpm and mRNA was isolated using Oligotex following the manufacturer's instructions. RT-PCT was performed using an Access RT-PCR kit (Promega). One μ g of total RNA was used in each reaction together with specific oligonucleotide primers (50 pmol) for the following genes as indicated under conditions according to the manufacturer's instructions. PCR was performed using a Hybaid machine, dynazyme II polymerase, 1.5mM Mg, 28-35 cycles at an annealing temperature between 56-63°C.

25

15

30

20

Results The results are illustrated in Figure 3 and reveal the following changes in transcript levels for naive splenic B cells

35

Delta1	IL-10 increased	Notch2	IL-10 increased
	LPS increased		LPS increased
Serrate1	IL-10 no change	Notch4	IL-10 increased
	LPS increased		LPS increased

40

25

45

Example 11 - Modulation of the expression of Notch receptors and ligands on murine bone marrow derived dendritic cells (DCs) in response to IL-10

30

50

CD11c+ DCs were MACS separated from the spleens of naive BALB/C mice and incubated at 37°C for 24 hours in medium alone or together with IL-10 (50ng/ml). Cells

55

5 were collected and mRNA was isolated using Oligotex following the manufacturer's
 instructions. RT/PCR was performed using an Access RT-PCR kit (Promega). One µg of
 10 total RNA was used in each reaction together with specific oligonucleotide primers (50
 5 pmol) for the following genes as indicated under conditions according to the
 manufacturer's instructions. PCR was performed using a Hybaid machine, dynazyme II
 polymerase, 1.5 mM Mg, 28-35 cycles at an annealing temperature between 56-63°C.

15 **Results** The results are illustrated in Figure 4 and reveal the following changes in
 transcript levels for bone marrow derived DCs

10	Mature versus immature DCs		
20	Delta1	increased	
	Notch2	decreased	
	Serrate1	increased	
	IL-10 stimulation IL-10 of mature DCs		
25	15	Delta1	decreased
		Serrate1	increased

30 In Figure 4 the results show bone marrow-derived DC cultivated in the presence of IL-10
 immature DCs (Lane 1), mature DCs (Lane 2) and DCs cultivated with 50 ng/ml IL-10 (Lane
 20 3). Hes1 transcription is measured as an index of Notch signalling.

35 References

- Morgenstern and Land (1990) *Nucleic Acids Res.* 18 :3587-3596.
- 40 25 Cosset *et al.* (1995) *J. Virol.* 69: 7430-7436.
- Coffin RS, *et al.* (1998) *Gene Therapy* 5: 718-722.
- Inaba K, *et al.* (1992) *J. Exp. Med.* 175: 1157-1167.
- Caux C, *et al.* (1992) *Nature* 360: 258-261.
- 45 Sallusto F and Lanzavecchia A (1994) *J. Exp. Med.* 179: 1109-1118.
- 30 Zhao *et al.* (1995) *J. Immunol* 155:3904-3911.
- Wilson and Hemmati-Brivanlou (1997) *Neuron* 18: 699-710.
- 50 Hemmati-Brivanlou and Melton (1997) *Cell* 88: 13-17.

- 5 Valenzuela *et al.* (1995) *J. Neurosci* 15: 6077-6084.
Sasai *et al.* (1994) *Cell* 79: 779-790.
Shimizu *et al.* (1999) *J. Biol. Chem.* 274(46): 32961-32969.
- 10 Iemura *et al.* (1998) *PNAS* 95: 9337-9342.
- 5 Altman *et al.* (1996) *Science* 274: 94-96.
Dkuz *et al.* (1997) *Cell* 90: 271-280.

15

20

25

30

35

40

45

50

55

Claims

5

10

15

20

25

30

35

40

45

50

55

CLAIMS

5

1. A method for producing a lymphocyte or antigen presenting cell (APC) having tolerance to an allergen or antigen which method comprises incubating a lymphocyte or APC
10 obtained from a human or animal patient with (i) a composition capable of upregulating expression of an endogenous Notch or Notch ligand in the lymphocyte and/or APC and (ii) the allergen or antigen.

15

2. A method according to claim 1 wherein the method comprises incubating a lymphocyte or APC obtained from a human or animal patient with an APC in presence of (i)
20 a composition capable of upregulating expression of an endogenous Notch or Notch ligand in the lymphocyte and/or APC and (ii) the allergen or antigen.

20

3. A method according to claim 1 for producing an APC capable of inducing in a T cell tolerance to an allergen or antigen which method comprises contacting an APC with (i) a
25 composition capable of upregulating expression of an endogenous Notch or Notch ligand in the APC and (ii) the allergen or antigen.

25

4. A method according to claim 1 or claim 2 for producing *ex vivo* a T cell having tolerance to an allergen or antigen which method comprises incubating a T cell obtained from
30 a human or animal patient with an antigen presenting cell (APC) in the presence of (i) a composition capable of upregulating expression of an endogenous Notch or Notch ligand in the APC and/or T cell and (ii) the allergen or antigen.

30

35

5. A method according to any one of claims 1 to 4 wherein the composition comprises a polypeptide selected from Noggin, Chordin, Follistatin, Xnr3, FGF and derivatives,
40 fragments, variants and homologues thereof, and immunosuppressive cytokines, or a combination thereof.

40

45

6. A method according to claim 5 wherein the immunosuppressive cytokine is selected from IL-4, IL-10, IL-13, TGF- β and FLT3.

50

55

- 5 7. A method according to any one of the preceding claims wherein the Notch ligand is selected from Serrate, Delta and homologues thereof.
- 10 8. A method according to any one of the preceding claims wherein the APC is a dendritic cell.
- 15 9. A method for producing a lymphocyte or APC having tolerance to an allergen or antigen which method comprises incubating a lymphocyte or APC obtained from a human or animal patient with a lymphocyte or APC produced by the method of any one of the preceding claims.
- 20 10. A method according to claim 9 for producing *ex vivo* a T cell having tolerance to an allergen or antigen which method comprises incubating a T cell obtained from a human or animal patient with a T cell produced by the method of any one of the preceding claims.
- 25 11. Use of a lymphocyte or APC produced by the method of any one of the preceding claims in suppressing an immune response in a mammal to the allergen or antigen.
- 30 12. Use of a composition capable of upregulating expression of an endogenous Notch or Notch ligand in an APC or lymphocyte in a method of producing regulatory lymphocytes capable of suppressing the activity of other lymphocytes.
- 35 13. Use according to claim 12 wherein the composition is as defined in claims 5 or 6.
- 40 14. Use according to claim 12 or 13 wherein the Notch ligand is selected from Serrate, Delta and homologues thereof.
- 45 15. Use according to any one of claims 12 to 14 wherein the APC is a dendritic cell.
- 50 16. A method of treating a patient suffering from a disease characterised by inappropriate lymphocyte activity which method comprises administering to the patient a lymphocyte produced by the method of any one of claims 1 to 8.
- 55

5

17. A method for producing a lymphocyte having tolerance to an allergen or antigen which method comprises incubating an APC produced by the method of claim 3 with the lymphocyte.

10

15

20

25

30

35

40

45

50

55




FIG. 1




FIG. 2

SUBSTITUTE SHEET (RULE 26)

BEST AVAILABLE COPY




FIG. 3




FIG. 4

SUBSTITUTE SHEET (RULE 26)

BEST AVAILABLE COPY